

SNUTP-98-031

revised, Aug., 1998, hep-th/9804093

Orbifolds of $AdS_7 \times S^4$ and Six Dimensional $(0, 1)$ SCFT

CHANGHYUN AHN^a, KYUNGHO OH^b AND RADU TATAR^c

^a *Dept. of Physics, Seoul National University, Seoul 151-742, Korea*

`chahn@spin.snu.ac.kr`

^b *Dept. of Mathematics, University of Missouri-St. Louis, St. Louis, MO 63121, USA*

`oh@arch.ums1.edu`

^c *Dept. of Physics, University of Miami, Coral Gables, FL 33146, USA*

`tatar@phyvax.ir.miami.edu`

ABSTRACT

We analyze the relation between the large N limit of 6 dimensional superconformal field theories with eight supercharges and M theory on orbifolds of $AdS_7 \times \mathbf{S}^4$. We use the known spectrum of Kaluza-Klein harmonics of supergravity on $AdS_7 \times \mathbf{S}^4$ and we take their orbifold projection to get information about the chiral primary operators of 6 dimensional SCFT which is realized on the worldvolume of M5 brane sitting at the orbifold singularities.

1 Introduction

Recently, the connection between anti-de Sitter (AdS) space and the dynamics of world-volume of D brane has been actively studied. Maldacena [1] proposed that the large N limit of super conformal field theories (SCFT) can be described by taking the supergravity limit of the superstring compactified on AdS space. Furthermore, the correlation functions in the SCFT that has the ADS boundary at infinity boundary as its spacetime can be obtained by taking the dependence of the supergravity action on the asymptotic behavior of its fields at the boundary [2, 3]. This way one can get the scaling dimensions of operators in the SCFT from the masses of particles in string/M theory.

Moreover, for any given $\mathcal{N} = 2, 1, 0$ superconformal model in 4 dimensions there exists a formulation of type IIB theory compactified on orbifolds of $AdS_5 \times \mathbf{S}^5$ [4, 5]. This proposed duality has been tested by studying the Kaluza-Klein (KK) states of theory on the orbifolds of $AdS_5 \times \mathbf{S}^5$ and by comparing them with the chiral primary operators of the SCFT on the boundary [6].

As we go one step further, the field theory/ M theory duality gives M theory on the AdS_4 or AdS_7 for some superconformal theory in 3 and 6 dimensions, respectively. The maximally supersymmetric theories in 3 and 6 dimensions have been studied recently [7, 8, 9, 10, 11]. The lower supersymmetric case was realized on the worldvolume of M theory branes sitting on the orbifold singularities [12] (See also [13]). Very recently, along the line of [6], the Kaluza-Klein states of M theory on the orbifolds of $AdS_4 \times \mathbf{S}^7$ were studied and compared with the chiral primary operators of the SCFT on the boundary [14].

In this paper, we investigate the Kaluza-Klein states of M theory for an orbifolds of $AdS_7 \times \mathbf{S}^4$ and we obtain the dimensions of operators in the superconformal multiplets by using the correspondence between AdS compactifications and SCFT. We obtain results referring only to the untwisted sectors of M theory compactified on AdS space which is the one that can be derived from supergravity. By taking all the degrees of freedom of M theory the singular geometry is regulated.

2 Review on AdS_7 and $(0, 2)$ SCFT

We briefly review the SCFT/ AdS_7 relation proposed in [2, 3]. The boundary of AdS_7 is a 6 dimensional Minkowski space with points at infinity added. The isometry group of AdS_7 is $SO(6, 2)$ which is also the conformal group on the boundary. The proposed

duality relates M theory on AdS_7 to the large N limit of SCFT on its boundary. In the Euclidean version the boundary is 6 sphere $\mathbf{S}^6$. Consider the maximally supersymmetric case, so that the internal space is also a sphere. Let ϕ be a scalar field on AdS_7 and ϕ_0 its restriction to the boundary $\mathbf{S}^6$. According to the SCFT/ AdS_7 correspondence, ϕ_0 couples to a conformal operator $\mathcal{O}$ on the boundary through a coupling $\int_{\mathbf{S}^6} \phi_0 \mathcal{O}$.

When ϕ has mass m , the corresponding operator $\mathcal{O}$ has the conformal dimension Δ given by the following relation

$$m^2 = \Delta(\Delta - 6). \quad (2.1)$$

The irrelevant, marginal and relevant operators of the boundary theory correspond to massive, massless and tachyonic modes in the supergravity theory. If a p -form C on AdS_7 couples to a $6-p$ form operator $\mathcal{O}$ on the boundary through a coupling $\int_{\mathbf{S}^6} C \wedge \mathcal{O}$, then the relation between the mass of C and the conformal dimension of $\mathcal{O}$ is given by

$$m^2 = (\Delta + p)(\Delta + p - 6). \quad (2.2)$$

The value of m^2 in this formula is the eigenvalue of the Laplace operator on the AdS_7 space in the supergravity side. The values usually quoted for p -forms are the eigenvalues $\widetilde{m}^2$ of the appropriate Maxwell-like operators. Their relation with the dimensions is given by [15, 7]

$$\widetilde{m}^2 = (\Delta - p)(\Delta + p - 6). \quad (2.3)$$

The massless graviton in the AdS_7 supergravity couples to the dimension 6 stress-energy tensor of the SCFT. When the internal space has continuous rotational symmetry, there exist also AdS_7 massless vector fields in its adjoint representation which can couple to the five dimensional R symmetry currents of the SCFT.

Let us consider M theory on $AdS_7 \times \mathbf{S}^4$ with a 4-form flux of N quanta on $\mathbf{S}^4$, and with the radii of the AdS_7 and $\mathbf{S}^4$ being $R_{AdS_7} = 2R_{S^4} = 2l_p(\pi N)^{1/3}$. The eleven dimensional supergravity is appropriate at energies much smaller than the Planck scale $1/l_p$. For large N this includes the energy range of the Kaluza-Klein modes, whose mass is of the order of $1/R_{AdS_7}$. The bosonic symmetry of this compactification of eleven dimensional supergravity is $SO(6, 2) \times SO(5)$. In [1] it was proposed that the $(0, 2)$ conformal theory, which is the decoupled theory on N parallel M5 branes, is dual to M theory on the above background. The $SO(6, 2)$ part of the symmetry of the supergravity theory is the conformal group of the SCFT, which can be thought of as living on the boundary of the AdS_7 space. The $SO(5)$ part of the symmetry corresponds to the R symmetry of the superconformal theory.

The Kaluza-Klein excitations of supergravity, in the maximally supersymmetric cases fall into small representations of supersymmetry. Thus, their mass formula is protected from quantum and string/M theory corrections. According to the proposal in [3], they

can couple to chiral fields of the SCFT on the boundary, for which the scaling dimensions are also protected from quantum corrections.

The spectrum of the Kaluza-Klein harmonics of M theory on $AdS_7 \times \mathbf{S}^4$ was analyzed in [16]. There are three families of scalar excitations. Two families contain states with only positive m^2 and which only correspond to irrelevant operators. One family has also states with negative and zero m^2 . They fall into the k -th order symmetric traceless representation of $SO(5)$ with unit multiplicity. Their masses are *

$$m^2 = 4k(k-3), \quad k = 2, 3, \dots \quad (2.4)$$

The field corresponding to $k = 1$ also appears in the supergravity, and this is the singleton which may be gauged away everywhere except at the boundary of the AdS space and decouples from all other operators. By using (2.1), the dimensions of the corresponding operators in the SCFT are

$$\Delta = 2k, \quad k = 2, 3, \dots \quad (2.5)$$

These are precisely the dimensions of the chiral primary operators found in [17], which parameterize the space of flat directions $(\mathbf{R}^5)^N/S_N$ in a gauge invariant way and this can be viewed as a test of the conjecture in [1].

There is one family of vector bosons that also contains massless states

$$\widetilde{m}^2 = 4(k^2 - 1), \quad k = 1, 2, \dots \quad (2.6)$$

By using (2.3), the dimensions of the corresponding 1-form operators in the SCFT are given by

$$\Delta = 2k + 3, \quad k = 1, 2, \dots \quad (2.7)$$

The massless vector obtained for $k = 1$ in (2.6) corresponds to the dimension 5 R symmetry current in SCFT. In general, chiral fields corresponding to all the towers of Kaluza-Klein harmonics are related to the scalar operators of (2.5) by the superconformal algebra, as discussed in the four dimensional case in [15]. Each value of k gives rise (at least for large enough k) to one field in each tower of KK states, with an $SO(5)$ representation that is determined by the representation of the scalar field. In particular, the R symmetry currents and the stress energy tensor sit in the same superconformal representation as the scalar field with $k = 2$ mentioned above.

3 An Orbifold of $AdS_7 \times S^4$ and $(0, 1)$ SCFT

It is proposed that the $(0, 1)$ SCFT on N M5 branes can be constructed by investigating M theory on the orbifold of $AdS_7 \times \mathbf{S}^4$. The result of [12] states that M theory on

*The masses should be scaled by a factor of $e = 2$.

$AdS_7 \times \mathbf{S}^4/\mathbf{Z}_k$ is dual to $(0,1)$ six dimensional theory with the gauge group $SU(k)^{N-1}$ and with one tensor multiplet and k hypermultiplets in the $\mathbf{k} \oplus \bar{\mathbf{k}}$ representation of the gauge group factors. The theory has a $SU(2)_R$ symmetry, the only bosons which are transforming non-trivially under this symmetry being the ones in the hypermultiplets and these transform in the fundamental representation. For the M5-brane, the transverse space is $\mathbf{R}^5$ and an appropriate orbifolding leads to $(0,1)$ theories on the world-volume of the M5-brane. In order to keep the AdS_7 structure in the near-horizon geometry, the orbifolding should act on $\mathbf{S}^4$ only [12].

One approach to obtain such orbifolds would be to break $\mathbf{S}^4$ into fiberations of smaller dimensional spheres over a base manifold. Then one can obtain an orbifold by taking actions on the base. The $SO(5)$ isometry of $\mathbf{S}^4$ will be broken into the isometry groups of both the fiber and the base. By embedding a discrete group into the isometry of the base corresponding to the orbifolding process, the isometry group of the base will be further broken. This method has been inspired by the work of Entin and Gomis [14] in the context of $AdS_4 \times \mathbf{S}^7$.

To explain this explicitly, let

$$\mathbf{S}^4 = \{(x_1, x_2, x_3, x_4, x_5) \in \mathbf{R}^5 \mid x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_5^2 = \rho\} \quad (3.1)$$

$$\mathbf{D}^i = \{(x_1, \dots, x_i) \in \mathbf{R}^i \mid x_1^2 + \dots + x_i^2 \leq \rho\} \quad (3.2)$$

where ρ is the radius of the 4 sphere $\mathbf{S}^4$ given by $\rho = R_{S^4} = l_p(\pi N)^{1/3}$ as we mentioned previous section. We list the possible fiberations of $\mathbf{S}^4$ induced by the coordinate projections.

- Case 1

$$\pi_1 : \mathbf{S}^4 \rightarrow \mathbf{D}^1, \quad (x_1, x_2, x_3, x_4, x_5) \rightarrow (x_1) \quad (3.3)$$

Then the generic fiber will be a 3 sphere $\mathbf{S}^3$ and the only isometry of the base leaving the boundary of the base invariant is $\mathbf{Z}_2$ whose action is given by $x_1 \leftrightarrow -x_1$. Now we take the orbifold projection by $\mathbf{Z}_2$. Then we obtain the hemisphere of Berkooz [13] which is made of a bundle of 3-sphere over a 1 dimensional manifold $\mathbf{D}^1/\mathbf{Z}_2$. Thus this theory has an $SO(4)$ symmetry. Furthermore, it is states in [13] that the field theory can acquire a global E_8 symmetry by placing E_8 vector gauge field on the fiber over $x_1 = 0$.

- Case 2

$$\pi_2 : \mathbf{S}^4 \rightarrow \mathbf{D}^2, \quad (x_1, x_2, x_3, x_4, x_5) \rightarrow (x_1, x_2) \quad (3.4)$$

The generic fiber is 2 sphere $\mathbf{S}^2$ and the the base $\mathbf{D}^2$ can be embedded into a complex z -plane $\mathbf{C}$ by letting $z = x_1 + ix_2$. Now the discrete subgroup $\mathbf{Z}_k$ of $U(1)$ can act on

the base $\mathbf{D}^2$. Under the $\mathbf{Z}_k$ projection, the $SO(3) = SU(2)$ symmetry is intact and only charges which are multiples of k are allowed in the theory.

- Case 3

$$\pi_3 : \mathbf{S}^4 \rightarrow \mathbf{D}^3, \quad (x_1, x_2, x_3, x_4, x_5) \rightarrow (x_1, x_2, x_3) \quad (3.5)$$

The generic fiber is a circle $\mathbf{S}^1$ and the discrete subgroups of $SU(2) = SO(3)$ can act on the base $\mathbf{D}^3$. Under the $\mathbf{Z}_k$ projection, the $U(1)$ symmetry coming from $\mathbf{S}^1$ is intact. We know that the $(0, 1)$ theory has a global symmetry $SU(2)_R$ and this cannot be realized because the $SU(2)$ symmetry is broken by the action of the $\mathbf{Z}_k$ group. So this case does not fit our goal.

- Case 4

$$\pi_4 : \mathbf{S}^4 \rightarrow \mathbf{D}^4, \quad (x_1, x_2, x_3, x_4, x_5) \rightarrow (x_1, x_2, x_3, x_4) \quad (3.6)$$

The generic fiber is two points (two roots for x_5) and the base can be embedded into a (u, v) -complex space $\mathbf{C}^2$ by letting $u = x_1 + ix_2, v = x_3 + ix_4$. Now we can break the isometry $SO(5)$ into $SO(4) \times \mathbf{Z}_2$. A discrete group Γ of ADE -type can act on the base $\mathbf{D}^4$. By embedding of Γ into the first factor $SU(2)_\Gamma$ of $SO(4) = SU(2)_\Gamma \times SU(2)_R$, $SO(5)$ has further broken into $\Gamma \times SU(2)_R$. This $\mathbf{Z}_2$ symmetry interchanges the upper hemisphere and the lower hemisphere.

In all four cases we have to be careful with the singular points. The orbifold acts geometrically on the bases and the origin of the disc is always a fixed point of the orbifold group. Supergravity alone cannot describe this fixed point. We would like to emphasize that we need to include all the degrees of freedom of M theory and not just the eleven dimensional supergravity ones in order to describe this fixed point. In addition to untwisted sectors, M theory includes twisted sectors and these will help to regulate the singular geometry at the origin of the disk. Another singularity appears on the boundary of the disc where the radius of the sphere fiber goes to zero. In this case, the fact that we use all the degrees of freedom of M theory again helps to regulate the singular geometry. Besides, the $\mathbf{S}^4$ radius ρ is proportional with N which implies that the disc radii are also proportional with N . So in the large N limit we can take the boundary arbitrarily far away. Therefore the origin and boundary singularities do not appear in the full M theory. In our paper we are obtaining the spectrum of chiral primary operators obtained from the untwisted sector of M theory. It should be very interesting to obtain the information coming from the twisted sectors of M theory.

We are going to work out the cases 2 and 4. First let us consider the case 4. We will exhibit the group actions Γ in details for the π_4 case:

$$A_{k-1}(\Gamma = \mathbf{Z}_k) : \quad u \rightarrow \exp\left(\frac{2\pi i}{k}\right) u, \quad v \rightarrow \exp\left(\frac{-2\pi i}{k}\right) v. \quad (3.7)$$

$$\begin{aligned}
D_{k-1}(\Gamma = \mathcal{D}_{k-2}) : \quad & u \rightarrow \exp\left(\frac{\pi i}{k-2}\right) u, \quad v \rightarrow \exp\left(\frac{-\pi i}{k-2}\right) v, \\
& u \rightarrow v, \quad v \rightarrow -u. \\
E_6(\Gamma = \mathcal{T}) : \quad & \text{binary tetrahedral group of order 24.} \\
E_7(\Gamma = \mathcal{O}) : \quad & \text{binary octahedral group of order 48.} \\
E_8(\Gamma = \mathcal{I}) : \quad & \text{binary icosahedral group of order 120.}
\end{aligned}$$

The Kaluza-Klein harmonics of M theory on $AdS_7 \times \mathbf{S}^4/\Gamma$ are Γ invariant states and can be obtained by Γ projection of the Kaluza-Klein harmonics on $AdS_7 \times \mathbf{S}^4$ discussed in previous section. Let us consider the scalar modes with mass formula given by (2.4) and check the relevant and marginal chiral primary operators in this family. We list $Sp(4) \rightarrow SU(2)_\Gamma \times SU(2)_R$ and $Sp(4) \rightarrow SU(2)_R$ branching rules for convenience in the Appendix A and B.

$$\bullet A_{k-1}(\Gamma = \mathbf{Z}_k)$$

Before discussing the surviving Kaluza-Klein modes, we briefly discuss the SCFT theory obtained on the M5 branes world-volume. As discuss in [12], we consider N M5 branes in a transverse space $\mathbf{R} \times \mathbf{C}^2/\mathbf{Z}_k$. In the type IIA side this representation is given by k D6 branes and N NS5 branes. The 6 dimensional world-volume theory is the $\mathcal{N} = 1, SU(k)^{N-1}$ gauge theory with one tensor multiplet and k hypermultiplets in the $\mathbf{k} \oplus \bar{\mathbf{k}}$ representation for each of the gauge group factors. Let us denote by $U_a^i, (i = 1, \dots, N-1; a = 1, \dots, k)$ the hypermultiplets. The field theory which is obtained is conformal. The $\mathcal{N} = 1$ theory in 6 dimensions has a $SU(2)_R$ R-symmetry. The scalars in the hypermultiplet transform as a $\mathbf{2}$ under $SU(2)_R$. The scalars in the vector multiplet transform as a $\mathbf{3}$ under $SU(2)_R$. We assign dimension 1 to all of them.

Now we proceed to discuss the surviving Kaluza-Klein modes.

We first note that $SU(2)_\Gamma$ representations can be realized as representations on the space of homogeneous polynomials in u, v and the invariant polynomials under the action of $\mathbf{Z}_k$ are generated by uv, u^k, v^k . Thus, for general k , we will keep only the odd dimensional representations of $SU(2)_\Gamma$ since these contain the powers of uv .

i) The $k = 1$ Kaluza-Klein particle in (2.4) transforms in the $\mathbf{5}$ of $SO(5)$. By decomposing $\mathbf{5}$ into representations of $SU(2)_\Gamma \times SU(2)_R$ [19, 20], we obtain

$$\mathbf{5} = (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}). \quad (3.8)$$

Now we perform the Γ projection on (3.8). The $(\mathbf{1}, \mathbf{1})$ is invariant under the Γ so the state $\mathbf{1}$ with respect to $SU(2)_R$ will survive the projection. However, these Kaluza-Klein modes can couple to dimension 2 operators. Only some of allowed projected states will be survived in the lower supersymmetry multiplets. A $(0, 1)$ superconformal subalgebra

of $(0, 2)$ superconformal algebra has a generator, R , of the $SU(2)_R$ symmetry. The dimensions of chiral operators are determined by $SU(2)_R$ symmetry half integer, so called spin s and by $SO(6)$ Lorentz quantum numbers. In this case $SO(6)$ weights are zero. Then the scaling dimensions of chiral primary operators are determined in terms of the spin as [18].

$$\Delta = 4s \quad (3.9)$$

This implies that a dimension 2 chiral primary operator has the spin $1/2$. Therefore **1** has the wrong spin to couple a dimension 2 chiral operator and we do not expect any dimension 2 chiral operators in the boundary $(0, 1)$ SCFT. Rather we would expect that the dimension 2 operators which are not chiral primary to couple to the Kaluza-Klein modes in the **1**. In the $(0, 2)$ SCFT, **(2, 2)** and **(1, 1)** in (3.8) sit in the same supermultiplet and the masses of **(1, 1)** Kaluza-Klein states were protected while for $(0, 1)$ SCFT, there is no such guarantee.

ii) The $k = 2$ Kaluza-Klein particle in (2.4) which transforms in the **14** of $SO(5)$ should couple to a dimension 4 chiral primary operator. Decomposing the **14** into representations of $SU(2)_\Gamma \times SU(2)_R$ gives

$$\mathbf{14} = (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}). \quad (3.10)$$

The **(3, 3)** and **(1, 1)** are invariants under the $\mathbf{Z}_k$ projection which give **3** and **1** under $SU(2)_R$. However, **3** is the only part which come from the decomposition (3.10) with the correct spin to couple a dimension 4 chiral primary operator according to (3.9). Therefore, we expect to have 3 dimension 4 chiral primary operators in the boundary SCFT. They are obtained by taking the gauge invariant operator $\text{Tr}(U^{i_1} U^{i_2} U^{i_3} U^{j_4})$ which is symmetric in the i 's indices and antisymmetric in j and either one of i 's. This is just in order to extract the **3** representation out of $\mathbf{2} \times \mathbf{2} \times \mathbf{2} \times \mathbf{2}$. The U^i are hypermultiplets transforming under consecutive gauge groups in order to obtain invariants.

iii) The $k = 3$ massless Kaluza-Klein particle in (2.4) should couple to dimension 6 chiral primary operators. It transforms in the **30** which decompose as

$$\mathbf{30} = (\mathbf{4}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}). \quad (3.11)$$

The **(3, 3)** and **(1, 1)** are invariant under the Γ projection. However, their spins are not consistent with coupling to a dimension 6 chiral operator from the analysis of (3.9). Therefore, there are no Kaluza-Klein harmonics in this family that can couple to dimension 6 chiral primary operators and no such operators are expected in the boundary $(0, 1)$ SCFT.

iv) The massless vector $k = 1$ in (2.6) is in the **10** of $SO(5)$. Decomposing **10** we get

$$\mathbf{10} = (\mathbf{3}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3}). \quad (3.12)$$

The $(\mathbf{3}, \mathbf{1})$ and $(\mathbf{1}, \mathbf{3})$ are invariant under the Γ projection. From the result of [18], isolated values of dimensions occur at $\Delta = 4s + 5$ for the vector. This implies that $(\mathbf{3}, \mathbf{1})$ is the only part in the decomposition (3.12) with the correct spin to couple to a dimension 5 chiral primary operator. Therefore, we expect the superconformal primary to be given by a $\mathbf{1}$ i.e. a singlet of $SU(2)_R$. In terms of the operator spectrum of the six dimensional SCFT a gauge invariant combination having dimension 5 and being a singlet of $SU(2)_R$ is $\text{Tr}(U\Phi UU^2)$ which is antisymmetric in the last two U 's.

It is rather straightforward to check the scaling dimensions with right quantum numbers for other massive representations: spinors, higher rank tensors and so on.

Now we turn to the binary dihedral group.

$$\bullet D_{k-1}(\Gamma = \mathcal{D}_{k-2}), E_n$$

Since the binary dihedral group $\mathcal{D}_{k-2}$ is generated by the cyclic group $\mathbf{Z}_{k-2/2}$ and a reflection, it will further restrict the representations. For our purpose, notice that $\mathbf{3}$ does not have an invariant component since uv is not invariant under the reflection. However, $\mathbf{1}$ has an invariant component for D_{k-1} . Again the 6 dimensional field theory brane configuration [12] is the same as the one for A_{k-1} gauge theory except that there exists an orientifold 6 plane. Each gauge group factors couple hypermultiplets in vector representation or fundamental representation. We expect the superconformal primary to be given by a $\mathbf{1}$ i.e. a singlet of $SU(2)_R$. In terms of the operator spectrum of the SCFT a gauge invariant combination having dimension 5 and being a singlet of $SU(2)_R$ is $\text{Tr}(U\Phi UU^2)$ which is antisymmetric in the last two U 's. Since $\mathcal{D}_2$ is the subgroup of E_6 and E_7 , $\mathbf{3}$ does not have an invariant component under these groups. Similarly, one can see that E_8 does not leave any invariant subspace of $\mathbf{3}$. It is true that $\mathbf{1}$ has an invariant component for E_n . But it is not clear how this can be realized in the SCFT side because there is no known field theory for E_n gauge theory.

Now we go to the case 2 i.e. to the projection π_2 as in (3.4). In this case the $SO(5)$ group is broken to $SU(2)_R \times U(1)/\mathbf{Z}_k$. The branching rule and (3.9) analysis tells us that there is no allowed Γ invariant states.

4 Conclusion

We studied the relation between chiral primary operators of SCFT in 6 dimensions and the KK states of M theory on orbifolds of $AdS_7 \times \mathbf{S}^4$. This generalizes the relation between the chiral primary operators of $(0, 2)$ SCFT and KK states of supergravity on

$AdS_7 \times \mathbf{S}^4$ found in [7, 8, 9, 10]. In our case, we have obtained the surviving KK modes in the orbifold models by projecting on Γ invariant states. the known KK states for supergravity compactified on $AdS_7 \times \mathbf{S}^4$.

Acknowledgments

We thank J. Gomis, M. Gunaydin, A. Kehagias, S. Minwalla, Y. Oz, J. Terning and A. Zaffaroni for email correspondences. RT would like to thank Prof. Orlando Alvarez for valuable discussions on various points in this paper. CA thanks B.-H. Lee for bring his attention to the ref. [19].

We would like to thank our referee for helping us to improve our paper.

5 Appendix A: $Sp(4)$ Branching Rule I

Fields	$Sp(4)$ Dynkin label	$SU(2)_\Gamma \times SU(2)_R$
$\sqrt{3}$ index anti. tensor	$(0, 0) : \mathbf{1}$	$(\mathbf{1}, \mathbf{1})$
spinor	$(1, 0) : \mathbf{4}$	$(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
scalar*	$(0, 1) : \mathbf{5}$	$(\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
graviton	$(0, 0) : \mathbf{1}$	$(\mathbf{1}, \mathbf{1})$
gravitino	$(1, 0) : \mathbf{4}$	$(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
$\sqrt{3}$ index anti. tensor	$(0, 1) : \mathbf{5}$	$(\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
vector*	$(2, 0) : \mathbf{10}$	$(\mathbf{3}, \mathbf{1})^\# \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$
scalar*	$(0, 2) : \mathbf{14}$	$(\mathbf{3}, \mathbf{3})^\# \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
spinor	$(1, 1) : \mathbf{16}$	$(\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
$\sqrt{3}$ index anti. tensor	$(0, 0) : \mathbf{1}$	$(\mathbf{1}, \mathbf{1})$
gravitino	$(1, 0) : \mathbf{4}$	$(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
graviton	$(0, 1) : \mathbf{5}$	$(\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
2 index anti. tensor	$(2, 0) : \mathbf{10}$	$(\mathbf{3}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$
$\sqrt{3}$ index anti. tensor	$(0, 2) : \mathbf{14}$	$(\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
gravitino	$(1, 1) : \mathbf{16}$	$(\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
spinor	$(3, 0) : \mathbf{20}$	$(\mathbf{4}, \mathbf{1}) \oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{4})$
scalar*	$(0, 3) : \mathbf{30}$	$(\mathbf{4}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
vector	$(2, 1) : \mathbf{35}$	$(\mathbf{4}, \mathbf{2}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{1})$ $\oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$
spinor	$(1, 2) : \mathbf{40}$	$(\mathbf{4}, \mathbf{3}) \oplus (\mathbf{3}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3})$ $\oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$

Table 1. The first 3 superconformal multiplets, their $Sp(4)$ Dynkin labelling and branching rules for $SU(2)_\Gamma \times SU(2)_R$. The asterisk * denote the spectrums of negative or zero m^2 . The two states with # are Γ invariant ones with appropriate right quantum number.

Fields	$Sp(4)$ Dynkin label	$SU(2)_\Gamma \times SU(2)_R$
scalar	$(0, 0) : \mathbf{1}$	$(\mathbf{1}, \mathbf{1})$
spinor	$(1, 0) : \mathbf{4}$	$(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
$\sqrt{3}$ index anti. tensor	$(0, 1) : \mathbf{5}$	$(\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
vector	$(2, 0) : \mathbf{10}$	$(\mathbf{3}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$
graviton	$(0, 2) : \mathbf{14}$	$(\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
gravitino	$(1, 1) : \mathbf{16}$	$(\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
spinor	$(3, 0) : \mathbf{20}$	$(\mathbf{4}, \mathbf{1}) \oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{4})$
$\sqrt{3}$ index anti. tensor	$(0, 3) : \mathbf{30}$	$(\mathbf{4}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1})$
2 index anti. tensor	$(2, 1) : \mathbf{35}$	$(\mathbf{4}, \mathbf{2}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{1})$ $\oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$
scalar	$(4, 0) : \mathbf{35}$	$(\mathbf{5}, \mathbf{1}) \oplus (\mathbf{4}, \mathbf{2}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{4})$ $\oplus (\mathbf{1}, \mathbf{5})$
gravitino	$(1, 2) : \mathbf{40}$	$(\mathbf{4}, \mathbf{3}) \oplus (\mathbf{3}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3})$ $\oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
scalar	$(0, 4) : \mathbf{55}$	$(\mathbf{5}, \mathbf{5}) \oplus (\mathbf{4}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{2})$ $\oplus (\mathbf{1}, \mathbf{1})$
spinor	$(3, 1) : \mathbf{64}$	$(\mathbf{5}, \mathbf{2}) \oplus (\mathbf{4}, \mathbf{3}) \oplus (\mathbf{3}, \mathbf{4}) \oplus (\mathbf{2}, \mathbf{5})$ $\oplus (\mathbf{4}, \mathbf{1}) \oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{4})$
spinor	$(1, 3) : \mathbf{80}$	$(\mathbf{5}, \mathbf{4}) \oplus (\mathbf{4}, \mathbf{5}) \oplus (\mathbf{4}, \mathbf{3}) \oplus (\mathbf{3}, \mathbf{4})$ $\oplus (\mathbf{3}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$
vector	$(2, 2) : \mathbf{81}$	$(\mathbf{5}, \mathbf{3}) \oplus (\mathbf{4}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{5}) \oplus (\mathbf{4}, \mathbf{2})$ $\oplus (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{2}, \mathbf{4}) \oplus (\mathbf{3}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$

Table 2. The fourth superconformal multiplet, its $Sp(4)$ Dynkin labelling and branching rules for $SU(2)_\Gamma \times SU(2)_R$.

6 Appendix B: $Sp(4)$ Branching Rule II

Fields	$Sp(4)$ Dynkin label	$SU(2)_R$
$\sqrt{3}$ index anti. tensor	$(0, 0) : \mathbf{1}$	$\mathbf{1}$
spinor	$(1, 0) : \mathbf{4}$	$\mathbf{4}$
scalar*	$(0, 1) : \mathbf{5}$	$\mathbf{5}$
graviton	$(0, 0) : \mathbf{1}$	$\mathbf{1}$
gravitino	$(1, 0) : \mathbf{4}$	$\mathbf{4}$
$\sqrt{3}$ index anti. tensor	$(0, 1) : \mathbf{5}$	$\mathbf{5}$
vector*	$(2, 0) : \mathbf{10}$	$\mathbf{7} \oplus \mathbf{3}$
scalar*	$(0, 2) : \mathbf{14}$	$\mathbf{9} \oplus \mathbf{5}$
spinor	$(1, 1) : \mathbf{16}$	$\mathbf{8} \oplus \mathbf{6} \oplus \mathbf{2}$
$\sqrt{3}$ index anti. tensor	$(0, 0) : \mathbf{1}$	$\mathbf{1}$
gravitino	$(1, 0) : \mathbf{4}$	$\mathbf{4}$
graviton	$(0, 1) : \mathbf{5}$	$\mathbf{5}$
2 index anti. tensor	$(2, 0) : \mathbf{10}$	$\mathbf{7} \oplus \mathbf{3}$
$\sqrt{3}$ index anti. tensor	$(0, 2) : \mathbf{14}$	$\mathbf{9} \oplus \mathbf{5}$
gravitino	$(1, 1) : \mathbf{16}$	$\mathbf{8} \oplus \mathbf{6} \oplus \mathbf{2}$
spinor	$(3, 0) : \mathbf{20}$	$\mathbf{10} \oplus \mathbf{6} \oplus \mathbf{4}$
scalar*	$(0, 3) : \mathbf{30}$	$\mathbf{13} \oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{1}$
vector	$(2, 1) : \mathbf{35}$	$\mathbf{11} \oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{5} \oplus \mathbf{3}$
spinor	$(1, 2) : \mathbf{40}$	$\mathbf{12} \oplus \mathbf{10} \oplus \mathbf{8} \oplus \mathbf{6} \oplus \mathbf{4}$

Table 3. The first 3 superconformal multiplets, their $Sp(4)$ Dynkin labelling and branching rules for $SU(2)_R$. The asterisk * denote the spectrums of negative or zero m^2 .

Fields	$Sp(4)$ Dynkin label	$SU(2)_R$
scalar	$(0, 0) : \mathbf{1}$	$\mathbf{1}$
spinor	$(1, 0) : \mathbf{4}$	$\mathbf{4}$
$\sqrt{3}$ index anti. tensor	$(0, 1) : \mathbf{5}$	$\mathbf{5}$
vector	$(2, 0) : \mathbf{10}$	$\mathbf{7} \oplus \mathbf{3}$
graviton	$(0, 2) : \mathbf{14}$	$\mathbf{9} \oplus \mathbf{5}$
gravitino	$(1, 1) : \mathbf{16}$	$\mathbf{8} \oplus \mathbf{6} \oplus \mathbf{2}$
spinor	$(3, 0) : \mathbf{20}$	$\mathbf{10} \oplus \mathbf{6} \oplus \mathbf{4}$
$\sqrt{3}$ index anti. tensor	$(0, 3) : \mathbf{30}$	$\mathbf{13} \oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{1}$
2 index anti. tensor	$(2, 1) : \mathbf{35}$	$\mathbf{11} \oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{5} \oplus \mathbf{3}$
scalar	$(4, 0) : \mathbf{35}$	$\mathbf{13} \oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{5} \oplus \mathbf{1}$
gravitino	$(1, 2) : \mathbf{40}$	$\mathbf{12} \oplus \mathbf{10} \oplus \mathbf{8} \oplus \mathbf{6} \oplus \mathbf{4}$
scalar	$(0, 4) : \mathbf{55}$	$\mathbf{17} \oplus \mathbf{13} \oplus \mathbf{11} \oplus \mathbf{9} \oplus \mathbf{5}$
spinor	$(3, 1) : \mathbf{64}$	$\mathbf{14} \oplus \mathbf{12} \oplus \mathbf{10} \oplus \mathbf{8}$ $\oplus \mathbf{8} \oplus \mathbf{6} \oplus \mathbf{4} \oplus \mathbf{2}$
spinor	$(1, 3) : \mathbf{80}$	$\mathbf{16} \oplus \mathbf{14} \oplus \mathbf{12} \oplus \mathbf{10}$ $\oplus \mathbf{10} \oplus \mathbf{8} \oplus \mathbf{6} \oplus \mathbf{4}$
vector	$(2, 2) : \mathbf{81}$	$\mathbf{15} \oplus \mathbf{13} \oplus \mathbf{11} \oplus \mathbf{11}$ $\oplus \mathbf{9} \oplus \mathbf{7} \oplus \mathbf{7} \oplus \mathbf{5} \oplus \mathbf{3}$

Table 4. The fourth superconformal multiplet, its $Sp(4)$ Dynkin labelling and branching rules for $SU(2)_R$.

References

- [1] J. Maldacena, hep-th/9711200.
- [2] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, hep-th/9802109.
- [3] E. Witten, hep-th/9802150.

- [4] S. Kachru and E. Silverstein, hep-th/9802183.
- [5] A. Lawrence, N. Nekrasov and C. Vafa, hep-th/9803015.
- [6] Y. Oz and J. Terning, hep-th/9803167.
- [7] O. Aharony, Y. Oz and Z. Yin, hep-th/9803051.
- [8] R.G. Leigh and M. Rozali, hep-th/9803068.
- [9] S. Minwalla, hep-th/9803053.
- [10] E. Halyo, hep-th/9803077.
- [11] J. Gomis, hep-th/9803119.
- [12] S. Ferrara, A. Kehagias, H. Partouche and A. Zaffaroni, hep-th/9803109.
- [13] M. Berkooz, hep-th/9802195.
- [14] R. Entin and J. Gomis, hep-th/9804060.
- [15] S. Ferrara, C. Fronsali and A. Zaffaroni, hep-th/9802203.
- [16] M. Gunaydin, P. van Nieuwenhuizen and N.P. Warner, Nucl. Phys. **B255** (1985) 63; P. van Nieuwenhuizen, Class. Quant. Grav. **2** (1985) 1.
- [17] O. Aharony, M. Berkooz and N. Seiberg, hep-th/9712117.
- [18] S. Minwalla, hep-th/9712074.
- [19] J. Patera and D. Sankoff, Tables of Branching rules for Representations of Simple Lie Algebras (L'Université de Montréal, Montréal, 1973).
- [20] W. MacKay and J. Patera, Tables of Dimensions, Indices and Branching Rules for representations of Simple Algebras (Dekker, New York, 1981).